


Numerical simulations of the Tayler-Spruit dynamo in proto-magnetars

Paul Barrère¹ *, Jérôme Guilet¹, Raphaël Raynaud² and Alexis Reboul-Salze³

¹Université Paris-Saclay, Université Paris Cité, CEA, CNRS, AIM, 91191, Gif-sur-Yvette, France

²Université Paris Cité, Université Paris-Saclay, CNRS, CEA, AIM, F-91191 Gif-sur-Yvette, France

³Max Planck Institute for Gravitational Physics (Albert Einstein Institute), 14476 Potsdam, Germany

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

The Tayler-Spruit dynamo is one of the most promising mechanisms proposed to explain angular momentum transport during stellar evolution. Its development in proto-neutron stars spun-up by supernova fallback has also been put forward as a scenario to explain the formation of very magnetized neutron stars called magnetars. Using three-dimensional direct numerical simulations, we model the proto-neutron star interior as a stably stratified spherical Couette flow with the outer sphere that rotates faster than the inner one. We report the existence of two subcritical dynamo branches driven by the Tayler instability. They differ by their equatorial symmetry (dipolar or hemispherical) and the magnetic field scaling, which is in agreement with different theoretical predictions (by Fuller and Spruit, respectively). The magnetic dipole of the dipolar branch is found to reach intensities compatible with observational constraints on magnetars.

Key words: stars: magnetars – supernovae: general – MHD – instabilities – magnetic fields

1 INTRODUCTION

Magnetars are a class of neutron stars that exhibit magnetic fields whose dipolar component reaches 10^{14} – 10^{15} G, which makes them the strongest fields observed in the Universe. Their dissipation are thought to power a wide variety of emissions like giant flares (Evans et al. 1980; Hurley et al. 1999, 2005; Svinikin et al. 2021), fast radio bursts (CHIME/FRB Collaboration et al. 2020; Bochenek et al. 2020), and short chaotic X-ray bursts (Gotz et al. 2006; Coti Zelati et al. 2018, 2021). Combined with a millisecond rotation, they may produce magnetorotational explosions, which are more energetic than standard supernovae explosions (Burrows et al. 2007; Dessart et al. 2008; Takiwaki et al. 2009; Kuroda et al. 2020; Bugli et al. 2020, 2021, 2023; Obergaulinger & Aloy 2020, 2021, 2022). The origin of these magnetic fields is therefore a crucial question to understand magnetars and their association to extreme events such as gamma-ray bursts or fast radio bursts. Two classes of scenarios can be distinguished for magnetar formation: (i) the merger of a neutron star binary, which may explain the plateau phase and the extended emission in X-ray sources associated with short gamma-ray bursts (Metzger et al. 2008; Lü & Zhang 2014; Gompertz et al. 2014). These events are interesting for their multimessenger signature but are expected to be too rare to be the main formation channel of Galactic magnetars, (ii) the core-collapse of a massive star, which is confirmed by the observation of Galactic magnetars associated with supernova remnants (Vink & Kuiper 2006; Martin et al. 2014; Zhou et al. 2019). In the latter case, the amplification of the magnetic field may be due either to the magnetic flux conservation during the collapse of the iron core of the progenitor star (Ferrario & Wickramasinghe 2006; Hu & Lou 2009; Schneider et al. 2020) or to a dynamo action in the newly born proto-magnetar. Indeed, two dynamo mechanisms

have already been studied by numerical simulations: the convective dynamo (Thompson & Duncan 1993; Raynaud et al. 2020, 2022; Masada et al. 2022; White et al. 2022) and the magnetorotational instability (MRI)-driven dynamo (Obergaulinger et al. 2009; Mösta et al. 2014; Reboul-Salze et al. 2021, 2022; Guilet et al. 2022). They have been shown to produce magnetar-like magnetic fields for millisecond rotation periods of the proto-magnetar, especially for periods $P \lesssim 10$ ms for the convective dynamo (Raynaud et al. 2020, 2022). These scenarios rely on the hypothesis that the rotation of the proto-magnetar is determined by the rotation of the progenitor core. However, it is still uncertain whether there are enough fast rotating progenitor cores to form all the observed magnetars in the Milky Way, which represent $\sim 10 - 40\%$ of the Galactic neutron star population (Kouveliotou et al. 1994; Woods & Thompson 2006; Gill & Heyl 2007; Beniamini et al. 2019).

In Barrère et al. (2022), we developed a new magnetar formation scenario in which the rapid rotation rate of the proto-magnetar is not determined by the progenitor core but by the ejected matter that remains gravitationally bound to the proto-magnetar and eventually falls back on the proto-magnetar surface $\sim 5 - 10$ s after the core-collapse. Since the accretion is asymmetric, the fallback matter transfers a significant amount of angular momentum to the surface (Chan et al. 2020; Janka et al. 2022), which makes the surface rotate faster than the core. In Barrère et al. (2022), we argue that this spin-up triggers the amplification of the magnetic field through the Tayler-Spruit dynamo mechanism. This dynamo mechanism can be described as a loop: (i) a poloidal magnetic field is sheared into a toroidal one (Ω -effect), (ii) the toroidal field becomes Tayler unstable after reaching a critical value (Tayler 1973; Pitts & Tayler 1985), and (iii) the Tayler instability regenerates a poloidal field (Fuller et al. 2019; Skoutnev et al. 2022; Ji et al. 2023).

The Tayler-Spruit dynamo was first modelled by Spruit (2002) to explain the angular momentum transport in stellar radiative zones.

* paul.barrere@cea.fr




Figure 1. Left: Bifurcation diagram of the time and volume averaged Elsasser number (and root mean square magnetic field) versus the Rossby number. Distinct dynamo branches are represented: dipolar (red), quadrupolar (mauve), hemispherical (green), and kinematic (black) whose respective thresholds are $Ro_D^c \sim 0.19$, $Ro_Q^c \sim 0.7$, $Ro_H^c \sim 0.37$, and $Ro_W^c \sim 0.62$. The hydrodynamic instability is triggered for $Ro_{\text{hyd}}^c > 0.177$. Dark green circles are stationary hemispherical dynamos and light green ones display parity modulations. Black crosses indicate failed dynamos, empty circles metastable solutions. Arrows attached to circles indicate the initial condition of the associated simulation. The black half empty circle specifies that the solution was found to be metastable in a simulation and stable in another. The error bars indicate the standard deviation. Right: snapshots of the magnetic field lines and surface radial fields associated to the different main dynamo branches at $Ro = 0.75$: dipolar (top), hemispherical (middle), and kinematic (bottom).

Fuller et al. (2019) provided a revised description, which tackles the previous critics of Spruit’s model (see Denissenkov & Pinsonneault 2007; Zahn et al. 2007) and yields different predictions. Their analytical results are now often included in stellar evolution codes (see e.g. Eggenberger et al. 2005; Cantiello et al. 2014; Eggenberger et al. 2019b,a; den Hartogh et al. 2020; Bonanno et al. 2020). Though this dynamo has long been debated in direct numerical simulations (Braithwaite 2006; Zahn et al. 2007), Petitdemange et al. (2023) recently reported a dynamo solution sharing many characteristics with the Tayler-Spruit model. Their numerical simulations modelled a stellar radiative zone, where the shear is negative, that is, the rotation rate decreases in the radial direction. In this Letter, we demonstrate that the Tayler instability can sustain different dynamo branches in the presence of positive shear, which gives strong support to the magnetar formation scenario of Barrère et al. (2022).

2 NUMERICAL SETUP

We perform three-dimensional (3D) direct numerical simulations of a stably stratified and electrically conducting Boussinesq fluid with the pseudo-spectral code MagIC (Wicht 2002; Gastine & Wicht 2012; Schaeffer 2013). The fluid has a constant density $\rho = 3.8 \times 10^{14} \text{ g cm}^{-3}$ (which corresponds to a proto-neutron star mass of $M = 1.4 M_{\odot}$) and evolves between two concentric spheres of radius r_i and $r_o = 12 \text{ km}$, rotating at the angular frequencies Ω_i and $\Omega_o = 2\pi \times 100 \text{ rad s}^{-1}$, respectively. The imposed differential rotation is characterized by the Rossby number $Ro \equiv 1 - \Omega_i/\Omega_o > 0$, which is varied between 0.125 and 1.2. This spherical Taylor-Couette configuration with positive shear prevents the development of the MRI and allows us to study the system in a statistically steady state. We impose no-slip and insulating boundary conditions at the inner and outer spheres. In all the simulations, we keep fixed

the other dimensionless control parameters: the shell aspect ratio $\chi \equiv r_i/r_o = 0.25$, the thermal and magnetic Prandtl numbers $Pr \equiv \nu/\kappa = 0.1$ and $Pm \equiv \nu/\eta = 1$, respectively, the Ekman number $E \equiv \nu/(d^2\Omega_o) = 10^{-5}$, and the ratio of the Brunt-Väisälä to the outer angular frequency $N/\Omega_o = 0.1$. The coefficients ν , κ , η , and $d \equiv r_o - r_i$ are respectively the kinematic viscosity, the thermal diffusivity, the resistivity, and the shell width. The magnetic energy is measured by the Elsasser number $\Lambda \equiv B_{\text{rms}}^2/(4\pi\rho\eta\Omega_o)$. The simulations are initialized either from a nearby saturated state, or with a weak ($\Lambda = 10^{-4}$) or a strong ($\Lambda = 10$) toroidal axisymmetric field with a given equatorial symmetry; it can be either dipolar (i.e. equatorially symmetric with $l = 2, m = 0$) or quadrupolar (i.e. anti-symmetric with $l = 1, m = 0$). We define a turbulent resistive time $\bar{\tau}_{\eta} = (\pi r_o/\bar{\ell})^2/\eta \sim 0.2d^2/\eta$, where $\bar{\ell} = 10$ is the typical value of the average harmonic degree of the time-averaged magnetic energy spectrum. In the following, we will term a solution *metastable* when a steady state is sustained for a time interval $\Delta t > 0.3\bar{\tau}_{\eta}$, and *stable* for $\Delta t \geq \bar{\tau}_{\eta}$ (up to $5.7\bar{\tau}_{\eta}$ for the simulation at $Ro = 0.2$).

3 RESULTS

We find in our set of simulations several dynamo branches represented by different colours in the bifurcation diagram shown in Fig. 1. When the differential rotation is low, the flow can not amplify a weak magnetic field (black crosses), but when $Ro > Ro_W^c \sim 0.62$, the magnetic field grows exponentially to reach a metastable or a stable saturated dynamo state (black dots). This kinematic dynamo is driven by a hydrodynamic instability of the Stewartson layer whose threshold is $Ro_{\text{hyd}}^c \sim 0.175$ (dashed vertical black line), which is in agreement with Hollerbach (2003). When $Ro \gtrsim 0.8$, the kinematic growth is followed by a non-linear growth and the system transitions directly to another branch with a larger magnetic energy (green cir-




Figure 2. Time-averaged ratio of the magnetic energy to the kinetic energy densities as a function of the Rossby number. The error bars indicate the standard deviation.

cles). Restarting from a nearby saturated solution or a strong toroidal field with quadrupolar symmetry (mauve dashed arrows), we find that the stability of this branch extends to Rossby number as low as $Ro_H^c \sim 0.37 < Ro_W^c$, which indicates that this dynamo is subcritical. By starting from a strong toroidal field with dipolar symmetry, we observe that this even subcritical branch is in bistability with another one which presents even stronger saturated magnetic fields $B_{rms} \in [4 \times 10^{14}, 1.1 \times 10^{15}]$ G (red circles). This branch is also subcritical since it can be maintained for Rossby numbers as low as $Ro > Ro_D^c \sim 0.19$. Moreover, the two subcritical branches do not only differ by their magnetic field strength but also by their equatorial symmetry, as seen in the 3D snapshots and the surface maps of the magnetic field in Fig. 1. Indeed, the magnetic field shows a dipolar symmetry on the stronger dynamo branch, whereas it is hemispherical on the weaker one. The latter can be interpreted as the superposition of modes with opposite equatorial symmetry (Gallet & Pétrelis 2009), which is consistent with the fact that we do find quadrupolar solutions (mauve circles in Fig. 1). These are only metastable for $Ro > Ro_Q^c \sim 0.7$ and transition to a stable dipolar or hemispherical solution. Finally, we note that the hemispherical dynamos with $Ro \geq 0.8$ (light green circles in Figs 1 and 2) display parity modulations (i.e. the solution evolves between hemispherical, dipole, and quadrupole symmetric states). This behaviour is reminiscent of the so-called Type 1 modulation identified in other dynamo setups (Knobloch et al. 1998; Raynaud & Tobias 2016) and likely results from the coupling of modes with opposite parity as the equatorial symmetry breaking of the flow increases at larger Rossby numbers.

The difference between the three dynamo branches is also clear in Fig. 2, where we see that the hemispherical branch saturates below the equipartition, with an energy ratio increasing with Ro from ~ 0.014 up to ~ 0.56 . By contrast, the dynamos of the dipolar branch are in a super-equipartition state ($E_b/E_k > 1$) and follow the magnetostrophic scaling $E_b/E_k \propto Ro^{-1}$ characteristic of the Coriolis-Lorenz force balance (Roberts & Soward 1972; Dormy 2016; Aubert et al. 2017; Dormy et al. 2018; Augustson et al. 2019; Seshasayanan & Gallet 2019; Raynaud et al. 2020; Schwaiger et al. 2019). This is also confirmed by force balance spectra shown in Fig. S1 in the Supplemental Materials.

Both subcritical dynamos show magnetic fields concentrated along the rotation axis, which differs significantly from the subcritical so-




Figure 3. Snapshots of the azimuthal slices of the angular velocity (left) and the magnetic field along the cylindrical radius $s \equiv r \sin \theta$ (right) of the dipolar dynamo at $Ro = 0.75$.

lutions found with a negative shear by (Petitdemange et al. 2023); this is also strikingly different from the magnetic field generated on the equatorial plane by the kinematic dynamo (see 3D snapshots of Fig. 1). This suggests that the dipolar and hemispherical dynamos are driven by a different mechanism. We argue that they are driven by the Tayler instability according to the following arguments. First, the axisymmetric toroidal magnetic component is clearly dominant since it contains 53 – 88 % of the total magnetic energy. Second, the simulations show a poloidal magnetic field with a dominant $m = 1$ mode (see Supplemental Materials Figs S2 and S3), which is the most unstable mode of the Tayler instability (Zahn et al. 2007; Ma & Fuller 2019). In the azimuthal cut of the magnetic field component B_s in Fig. 3, the Tayler mode also appears clearly close to the poles, where it is expected to develop for a toroidal field generated by the shearing of a poloidal field (Goossens et al. 1981). This is also consistent with the 3D snapshots of the dipolar and hemispherical branches in Fig. 1 where the toroidal magnetic field seems prone to a kink instability. Third, as in Petitdemange et al. (2023), the system bifurcates from the kinematic to the hemispherical branch in the vicinity of the threshold of the Tayler instability (Spruit 1999, 2002)

$$\Lambda_\phi^c \equiv \frac{B_\phi^2}{4\pi\rho\eta\Omega_o} \sim \frac{\chi}{1-\chi} \frac{N}{\Omega_o} \sqrt{\frac{Pr}{E}} \sim 3.3. \quad (1)$$

Indeed, if we focus on the stable and metastable kinematic solutions found at $Ro = 0.75$, we see in Fig. 4 that the *local* maximum of the toroidal axisymmetric field is in both cases close to the critical value above which it is expected to become unstable. The bifurcation from the kinematic toward the hemispherical branch that is observed for the metastable solution appears hence as the result of turbulent fluctuations departing far enough above the threshold of the Tayler instability.

Finally, we compare our numerical results to the theoretical predictions regarding the saturation of the Tayler-Spruit dynamo. Note that these predictions assume the scale separation $\omega_A \ll \Omega_o \ll N$, where the Alfvén frequency is defined by $\omega_A \equiv B_\phi / \sqrt{4\pi\rho r_o^2} \sim 12.1 (B_\phi / 10^{15} \text{ G}) \text{ Hz}$. Our numerical models assume $N/\Omega_o = 0.1$ to limit the computational costs, whereas for a typical PNS spun up by fallback to a period of 1 – 10 ms we expect $N/\Omega_o \sim 1 - 10$. On the other hand, the achieved magnetic field follows the right scale separation with $\omega_A/\Omega_o \lesssim 0.02$, which is expected to determine the




Figure 4. Time series of the maximum along the cylindrical radius s of the axisymmetric toroidal magnetic energy measured locally at $z = 0.45r_o$, for stable (black) and metastable (green) kinematic dynamos at $Ro = 0.75$. The dashed red line indicates the analytical threshold of the Taylor instability (equation 1). Dark lines show a running average and dotted green lines around $t \sim 20$ s indicate missing data.

saturation mechanism of the Taylor instability (Ji et al. 2023). Figure 5 displays the axisymmetric toroidal and poloidal magnetic fields (top), the dipole field (middle) and the Maxwell torque (bottom) as a function of an effective shear rate q measured locally in the saturated state of the dynamo (see Supplemental Materials Fig. S4). For the dipolar branch (red), we find that the power laws $B_{\text{tor}}^{m=0} \propto q^{0.36 \pm 0.05}$ and $B_{\text{pol}}^{m=0} \propto q^{0.62 \pm 0.07}$ fit the saturated magnetic field, while we find $B_s B_\phi \propto q^{1.0 \pm 0.02}$ or $B_s^{m=0} B_\phi^{m=0} \propto q^{1.1 \pm 0.04}$, depending on whether we take into account non-axisymmetric contributions to compute the Maxwell torque T_M . The scaling exponents are thus in good agreement with the theoretical predictions of Fuller et al. (2019) $B_{\text{tor}}^{m=0} \propto q^{1/3}$, $B_{\text{pol}}^{m=0} \propto q^{2/3}$ and $T_M \propto q$ (red dotted lines in Fig. 5). Contrary to their prediction, however, our torque is not dominated by the axisymmetric magnetic field, which may be related to their assumption of a stronger stratification. Interestingly, the hemispherical branch (green) does not follow the same scalings: for $q > 0.2$, we find $B_{\text{tor}}^{m=0} \propto q^{2.1 \pm 0.31}$ and $B_{\text{pol}}^{m=0} \propto q^{2.0 \pm 0.28}$ for the magnetic field, and $B_s B_\phi \propto q^{2.7 \pm 0.40}$ or $B_s^{m=0} B_\phi^{m=0} \propto q^{3.8 \pm 0.70}$ for the Maxwell torque. These results globally support Spruit’s predictions (Spruit 2002) $B_{\text{tor}}^{m=0} \propto q$, $B_{\text{pol}}^{m=0} \propto q^2$ and $T_M \propto q^3$ (green dotted lines), despite some tension concerning the scaling of the toroidal magnetic field. If we focus on the dipole field, we find the following power laws: $B_{\text{dip}} \propto q^{0.66 \pm 0.03}$ and $B_{\text{dip}} \propto q^{1.1 \pm 0.4}$, for the dipolar and hemispherical branches, respectively. The dipole field on the strong branch therefore follows the same scaling as the axisymmetric poloidal field and is only $\sim 33\%$ weaker.

4 CONCLUSIONS

To conclude, we show that the Taylor-Spruit dynamo also exists in the presence of positive shear. We demonstrate for the first time the existence of two subcritical branches of this dynamo with distinct equatorial symmetries, dipolar and hemispherical. Moreover, the former follows Fuller’s theoretical predictions, while the latter




Figure 5. Root mean square (RMS) toroidal and poloidal axisymmetric magnetic fields (top), RMS magnetic dipole (middle), and RMS magnetic torque (bottom) as a function of the time-averaged shear rate measured in the steady state, for the dipolar (red) and hemispherical (green) dynamo branches. Dotted lines show the best fits obtained with Fuller’s (red) and Spruit’s (green) theoretical scaling laws, respectively.

is in overall agreement with Spruit’s model. Compared to the study of Petitdemange et al. (2023) that use a negative shear, our results present a similar dynamical structure, with a bifurcation diagram characterized by a bistability between kinematic and subcritical dynamo solutions. The magnetic field of their Taylor-Spruit dynamo is, however, different since it is characterized by a smaller scale structure localized near the inner boundary in the equatorial plane, and induces a torque scaling according to Spruit’s prediction. Our study shows a magnetic field geometry concentrated near the pole in agreement with the expectation of the Taylor-Spruit dynamo and a more complex physics, with the existence of two different branches that can not be captured by a single scaling law. Extended parameter studies will be needed to further assess the impact of the resistivity and the stratification on this dynamo instability and better constrain its astrophysical implications.

Our results are of particular importance for stellar evolution models by confirming the existence of the Taylor-Spruit dynamo and by deepening our physical understanding of its complex dynamics. They also give strong support to the new magnetar formation scenario proposed by Barrère et al. (2022), which relies on the development of a Taylor-instability driven dynamo in the presence of a positive shear. We validate the assumption that the magnetic dipole is a significant

fraction of the poloidal magnetic field and follows the same scaling. Extrapolating our results for the dipolar branch to $q \sim 1$ as expected in Barrère et al. (2022), we obtain a magnetic dipole intensity of $\sim 3.2 \times 10^{14}$ G and an even stronger axisymmetric toroidal field of $\sim 2.1 \times 10^{15}$ G. These orders of magnitude are similar to those found in Barrère et al. (2022) for the same rotation period of $P_o \equiv 2\pi/\Omega_o = 10$ ms, and fall right in the magnetar range (Olausen & Kaspi 2014).

ACKNOWLEDGEMENTS

We thank F. Daniel, C. Gissinger and L. Petitdemange and F. Marcotte for fruitful discussions and for sharing the manuscript Petitdemange et al. (2023) before its publication. We also thank A. Igoshev for valuable comments. This work was supported by the European Research Council (MagBURST grant 715368), and the PNPS and PNHE programs of CNRS/INSU, co-funded by CEA and CNES. Numerical simulations have been carried out at the CINES on the Jean-Zay supercomputer (DARI project A130410317).

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.


REFERENCES

- Aubert J., Gastine T., Fournier A., 2017, *Journal of Fluid Mechanics*, 813, 558
- Augustson K. C., Brun A. S., Toomre J., 2019, *Astrophys. J.*, 876, 83
- Barrère P., Guilet J., Reboul-Salze A., Raynaud R., Janka H. T., 2022, *Astron. Astrophys.*, 668, A79
- Beniamini P., Hotokezaka K., van der Horst A., Kouveliotou C., 2019, *Mon. Not. Astron. R. Soc.*, 487, 1426
- Bochenek C. D., Ravi V., Belov K. V., Hallinan G., Kocz J., Kulkarni S. R., McKenna D. L., 2020, *Nature*, 587, 59
- Bonanno A., Guerrero G., Del Sordo F., 2020, *MEMSAI*, 91, 249
- Braithwaite J., 2006, *Astron. Astrophys.*, 449, 451
- Bugli M., Guilet J., Obergaulinger M., Cerdá-Durán P., Aloy M. A., 2020, *Mon. Not. Astron. R. Soc.*, 492, 58
- Bugli M., Guilet J., Obergaulinger M., 2021, *Mon. Not. Astron. R. Soc.*, 507, 443
- Bugli M., Guilet J., Foglizzo T., Obergaulinger M., 2023, *Mon. Not. Astron. R. Soc.*, 520, 5622
- Burrows A., Dessart L., Livne E., Ott C. D., Murphy J., 2007, *Astrophys. J.*, 664, 416
- CHIME/FRB Collaboration et al., 2020, *Nature*, 587, 54
- Cantiello M., Mankovich C., Bildsten L., Christensen-Dalsgaard J., Paxton B., 2014, *Astrophys. J.*, 788, 93
- Chan C., Müller B., Heger A., 2020, *Mon. Not. Astron. R. Soc.*, 495, 3751
- Coti Zelati F., Rea N., Pons J. A., Campana S., Esposito P., 2018, *Mon. Not. Astron. R. Soc.*, 474, 961
- Coti Zelati F., et al., 2021, *Astrophys. J. Lett.*, 907, L34
- Denissenkov P. A., Pinsonneault M., 2007, *Astrophys. J.*, 655, 1157
- Dessart L., Burrows A., Livne E., Ott C. D., 2008, *Astrophys. J. Lett.*, 673, L43
- Dormy E., 2016, *J. Fluid Mech.*, 789, 500
- Dormy E., Oruba L., Petitdemange L., 2018, *Fluid Dyn. Res.*, 50, 011415
- Eggenberger P., Maeder A., Meynet G., 2005, *Astron. Astrophys.*, 440, L9
- Eggenberger P., Buldgen G., Salmon S. J. A. J., 2019a, *Astron. Astrophys.*, 626, L1
- Eggenberger P., den Hartogh J. W., Buldgen G., Meynet G., Salmon S. J. A. J., Deheuvels S., 2019b, *Astron. Astrophys.*, 631, L6
- Evans W. D., et al., 1980, *Astrophys. J. Lett.*, 237, L7
- Ferrario L., Wickramasinghe D., 2006, *Mon. Not. Astron. R. Soc.*, 367, 1323
- Fuller J., Piro A. L., Jermyn A. S., 2019, *Mon. Not. Astron. R. Soc.*, 485, 3661
- Gallet B., Pétrélis F., 2009, *Phys. Rev. E*, 80, 035302
- Gastine T., Wicht J., 2012, *Icarus*, 219, 428
- Gill R., Heyl J., 2007, *Mon. Not. Astron. R. Soc.*, 381, 52
- Gompertz B. P., O'Brien P. T., Wynn G. A., 2014, *Mon. Not. Astron. R. Soc.*, 438, 240
- Goossens M., Biront D., Tayler R. J., 1981, *Astrophys. Space Sci.*, 75, 521
- Gotz D., et al., 2006, *The Astronomer's Telegram*, 953, 1
- Guilet J., Reboul-Salze A., Raynaud R., Bugli M., Gallet B., 2022, *Mon. Not. Astron. R. Soc.*, 516, 4346
- Hollerbach R., 2003, *J. Fluid Mech.*, 492, 289
- Hu R.-Y., Lou Y.-Q., 2009, *Mon. Not. Astron. R. Soc.*, 396, 878
- Hurley K., et al., 1999, *Nature*, 397, 41
- Hurley K., et al., 2005, *Nature*, 434, 1098
- Janka H.-T., Wongwathanarat A., Kramer M., 2022, *Astrophys. J.*, 926, 9
- Ji S., Fuller J., Lecoanet D., 2023, *Mon. Not. Astron. R. Soc.*, 521, 5372
- Knobloch E., Tobias S. M., Weiss N. O., 1998, *Mon. Not. Astron. R. Soc.*, 297, 1123
- Kouveliotou C., et al., 1994, *Nature*, 368, 125
- Kuroda T., Arcones A., Takiwaki T., Kotake K., 2020, *Astrophys. J.*, 896, 102
- Lü H.-J., Zhang B., 2014, *Astrophys. J.*, 785, 74
- Ma L., Fuller J., 2019, *Mon. Not. Astron. R. Soc.*, 488, 4338
- Martin J., Rea N., Torres D. F., Papitto A., 2014, *Mon. Not. Astron. R. Soc.*, 444, 2910
- Masada Y., Takiwaki T., Kotake K., 2022, *Astrophys. J.*, 924, 75
- Metzger B. D., Quataert E., Thompson T. A., 2008, *Mon. Not. Astron. R. Soc.*, 385, 1455
- Mösta P., et al., 2014, *Astrophys. J. Lett.*, 785, L29
- Obergaulinger M., Aloy M. Á., 2020, *Mon. Not. Astron. R. Soc.*, 492, 4613
- Obergaulinger M., Aloy M. Á., 2021, *Mon. Not. Astron. R. Soc.*, 503, 4942
- Obergaulinger M., Aloy M. Á., 2022, *Mon. Not. Astron. R. Soc.*, 512, 2489
- Obergaulinger M., Cerdá-Durán P., Müller E., Aloy M. A., 2009, *Astron. Astrophys.*, 498, 241
- Olausen S. A., Kaspi V. M., 2014, *Astrophys. J.*, 212, 6
- Petitdemange L., Marcotte F., Gissinger C., 2023, *Science*, 379, 300
- Pitts E., Tayler R. J., 1985, *Mon. Not. Astron. R. Soc.*, 216, 139
- Raynaud R., Tobias S. M., 2016, *J. Fluid Mech.*, 799, R6
- Raynaud R., Guilet J., Janka H.-T., Gastine T., 2020, *Sci. Adv.*, 6, eaay2732
- Raynaud R., Cerdá-Durán P., Guilet J., 2022, *Mon. Not. Astron. R. Soc.*, 509, 3410
- Reboul-Salze A., Guilet J., Raynaud R., Bugli M., 2021, *Astron. Astrophys.*, 645, A109
- Reboul-Salze A., Guilet J., Raynaud R., Bugli M., 2022, *Astron. Astrophys.*, 667, A94
- Roberts P. H., Soward A. M., 1972, *Annual Review of Fluid Mechanics*, 4, 117
- Schaeffer N., 2013, *Geochemistry, Geophysics, Geosystems*, 14, 751
- Schneider F. R. N., Ohlmann S. T., Podsiadlowski P., Röpke F. K., Balbus S. A., Pakmor R., 2020, *Mon. Not. Astron. R. Soc.*, 495, 2796
- Schwaiger T., Gastine T., Aubert J., 2019, *Geophysical Journal International*, 219, S101
- Seshasayanan K., Gallet B., 2019, *J. Fluid Mech.*, 864, 971
- Skoutnev V., Squire J., Bhattacharjee A., 2022, *Mon. Not. Astron. R. Soc.*, 517, 526
- Spruit H. C., 1999, *Astron. Astrophys.*, 349, 189
- Spruit H. C., 2002, *Astron. Astrophys.*, 381, 923
- Svinkin D., et al., 2021, *Nature*, 589, 211
- Takiwaki T., Kotake K., Sato K., 2009, *Astrophys. J.*, 691, 1360
- Tayler R. J., 1973, *Mon. Not. Astron. R. Soc.*, 161, 365
- Thompson C., Duncan R. C., 1993, *Astrophys. J.*, 408, 194
- Vink J., Kuiper L., 2006, *Mon. Not. Astron. R. Soc.*, 370, L14
- White C. J., Burrows A., Coleman M. S. B., Vartanyan D., 2022, *Astrophys. J.*, 926, 111
- Wicht J., 2002, *Physics of the Earth and Planetary Interiors*, 132, 281

- Woods P. M., Thompson C., 2006, in , Vol. 39, Compact stellar X-ray sources. pp 547–586, [doi:10.48550/arXiv.astro-ph/0406133](https://doi.org/10.48550/arXiv.astro-ph/0406133)
- Zahn J. P., Brun A. S., Mathis S., 2007, *Astron. Astrophys.*, 474, 145
- Zhou P., Vink J., Safi-Harb S., Miceli M., 2019, *Astron. Astrophys.*, 629, A51
- den Hartogh J. W., Eggenberger P., Deheuvels S., 2020, *Astron. Astrophys.*, 634, L16

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.

Supplementary materials for “Numerical simulations of the Tayler-Spruit dynamo in proto-magnetars”

Paul Barrère¹ *, Jérôme Guilet¹, Raphaël Raynaud² and Alexis Reboul-Salze³

¹Université Paris-Saclay, Université Paris Cité, CEA, CNRS, AIM, 91191, Gif-sur-Yvette, France

²Université Paris Cité, Université Paris-Saclay, CNRS, CEA, AIM, F-91191 Gif-sur-Yvette, France

³Max Planck Institute for Gravitational Physics (Albert Einstein Institute), 14476 Potsdam, Germany

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

In this Supplemental Material, we first detail the Boussinesq equations and numerical setup in Section 1 and give an overview of all simulations in Section 2. Then, we present the typical time-averaged spectra of the dynamos (Sections 3 and 4) and the method used to measure the local shear rate q in our simulations (Section 5).

1 GOVERNING EQUATIONS AND NUMERICAL SETUP

We model the proto-neutron star as a spherical stably stratified Couette flow whose surface rotates at the angular frequency $\Omega_o = 200\pi \text{ rad s}^{-1}$. In the rotating frame, the MHD equations under the Boussinesq approximation read

$$\nabla \cdot \mathbf{v} = 0, \quad (1)$$

$$D_t \mathbf{v} = -\frac{1}{\rho} \nabla p' - 2\Omega_o \mathbf{e}_z \times \mathbf{v} + \alpha g T' \mathbf{e}_r + \frac{1}{4\pi\rho} (\nabla \times \mathbf{B}) \times \mathbf{B} + \nu \Delta \mathbf{v}, \quad (2)$$

$$D_t T' = \kappa \Delta T', \quad (3)$$

$$\partial_t \mathbf{B} = \nabla \times (\mathbf{u} \times \mathbf{B}) + \eta \Delta \mathbf{B}, \quad (4)$$

$$\nabla \cdot \mathbf{B} = 0, \quad (5)$$

where \mathbf{v} is the velocity field, \mathbf{B} is the magnetic field, p' is the pressure perturbation, T' is the super-adiabatic temperature, ρ is the uniform density, $g = g_o r / r_o$ is the gravitation field, and $\alpha \equiv \rho^{-1} (\partial_T \rho)_p$ is the thermal expansivity. We assume that the viscosity ν , the thermal diffusivity κ and the magnetic diffusivity η are constant. We apply no-slip, electrically insulating, and fixed temperature boundary conditions on both shells. In the MagIC code, the length is scaled in units of shell thickness $d \equiv r_o - r_i$ where r_i and r_o are the inner and outer radii, the time in units of viscous time d^2/ν , the temperature in units of the fixed temperature gap between the shells $\Delta T \equiv T_o - T_i > 0$, and the magnetic field in units of $\sqrt{4\pi\rho\eta\Omega_o}$. The stratification is measured by the Rayleigh number $Ra \equiv N^2 d^4 / (\nu\kappa)$, where $N \equiv \sqrt{\alpha g_o \Delta T / d}$ is the Brunt-Väisälä frequency at the outer radius.

2 LIST OF SIMULATION MODELS

Tables 1 and 2 summarize the key parameters of the simulations carried out in this study.

arXiv:2306.12296v1 [astro-ph.HE] 21 Jun 2023

Table 1. Overview of the stable (or failed) dynamos solutions. The simulations are named using the value of the Rossby number followed by a letter which characterises the initial condition: ‘s’ for a strong dipole-like magnetic field (either analytic or from a saturated state of the dipolar branch), ‘as’ for a strong quadrupole-like field (analytic or from a saturated state of the quadrupolar branch), ‘h’ for the saturated state of the hemispherical branch, ‘k’ for a saturated state of the kinematic dynamo, and ‘w’ for a very weak seed field. The second column indicates the input Rossby number. The third column refers to the initial Elsasser number. The last column specifies the type of the dynamo solution: dipolar (DIP), quadrupolar (QUAD), hemispherical (HEM), oscillating (OSC), kinematic (WEAK), non dynamo (NONE). All the simulations have the same aspect ratio $\chi = 0.25$, Ekman number $E = 10^{-5}$, stratification $N/\Omega_o = 0.1$, thermal and magnetic Prandtl numbers $Pr = 0.1$ and $Pm = 1$, and the same resolution $(n_r, n_\theta, n_\phi) = (256, 256, 512)$.

Name	Ro	Λ_i	q	Λ	E_b/E_k	$B_{\text{tor}}^{m=0}$ [10^{14} G]	$B_{\text{pol}}^{m=0}$ [10^{14} G]	B_{dip} [10^{14} G]	$B_s B_\phi / 4\pi$ [10^{27} G 2]	$B_r B_\phi / 4\pi$ [10^{27} G 2]	$B_s^{m=0} B_\phi^{m=0} / 4\pi$ [10^{27} G 2]	$B_r^{m=0} B_\phi^{m=0} / 4\pi$ [10^{27} G 2]	Dynamo
Ro1.2s	1.2	$\Lambda(\text{Ro1s})$	0.13	70.11	1.73	9.16	1.04	0.70	41.97	45.70	8.31	25.11	DIP
Ro1s	1	$\Lambda(\text{Ro0.75s})$	0.069	87.83	4.74	10.46	1.17	0.77	49.37	53.33	10.80	30.60	DIP
Ro0.75s	0.75	10	0.06	51.85	3.93	8.08	0.68	0.48	22.57	21.10	4.43	10.24	DIP
Ro0.5s	0.5	$\Lambda(\text{Ro0.75s})$	0.032	33.06	6.16	6.65	0.46	0.33	11.31	9.84	2.20	4.02	DIP
Ro0.35s	0.35	$\Lambda(\text{Ro0.25s})$	0.02	20.87	8.36	5.34	0.39	0.25	7.01	6.39	1.53	2.80	DIP
Ro0.2s	0.2	$\Lambda(\text{Ro0.25s})$	0.0089	10.16	11.67	3.70	0.23	0.15	3.22	2.91	0.56	1.27	DIP
Ro0.17s	0.17	$\Lambda(\text{Ro0.25s})$	–	–	–	–	–	–	–	–	–	–	NONE
Ro0.125s	0.125	$\Lambda(\text{Ro0.25s})$	–	–	–	–	–	–	–	–	–	–	NONE
Ro1as	1	$\Lambda(\text{Ro0.75as})$	0.45	13.75	0.19	3.83	0.33	0.32	9.28	7.64	1.68	2.66	HEM
Ro0.75as	0.75	10	0.064	58.22	4.46	8.71	0.67	0.58	21.77	20.47	4.18	10.01	DIP
Ro0.65as	0.65	$\Lambda(\text{Ro0.75as})$	0.27	5.15	0.13	2.34	0.19	0.19	3.94	3.13	0.62	0.93	HEM
Ro0.5as	0.5	$\Lambda(\text{Ro0.75as})$	0.2	2.32	0.09	1.50	0.11	0.17	2.49	1.73	0.31	0.33	HEM
Ro0.4as	0.4	$\Lambda(\text{Ro0.75as})$	0.16	0.28	0.01	0.53	0.02	0.04	0.35	0.27	0.02	0.04	HEM
Ro1.2h	1.2	$\Lambda(\text{Ro1k})$	0.51	41.35	0.45	7.28	0.56	0.43	18.27	16.40	3.98	6.90	OSC
Ro0.42h	0.42	$\Lambda(\text{Ro0.5k})$	0.17	0.49	0.02	0.63	0.03	0.09	0.89	0.70	0.06	0.07	HEM
Ro1k	1	$\Lambda(\text{Ro0.75w})$	0.43	16.68	0.24	4.37	0.34	–	8.71	7.30	1.45	2.62	OSC
Ro0.5k	0.5	$\Lambda(\text{Ro0.75w})$	0.2	1.46	0.06	1.06	0.08	0.14	1.96	1.35	0.19	0.22	HEM
Ro0.85w	0.85	10^{-4}	0.33	14.32	0.26	4.07	0.31	0.36	7.58	6.56	1.41	2.57	OSC
Ro0.75w	0.75	10^{-4}	0.35	5.12	0.11	2.13	0.20	0.24	4.59	3.50	0.65	1.00	HEM
Ro0.75w2	0.75	10^{-4}	0.34	0.37	0.01	0.52	0.03	0.08	1.44	1.26	0.06	0.06	KIN
Ro0.65w	0.65	10^{-4}	0.27	4.87	0.13	2.16	0.18	–	3.93	3.11	0.60	0.87	HEM
Ro0.63w	0.63	10^{-4}	–	–	–	–	–	–	–	–	–	–	NONE
Ro0.6w	0.6	10^{-4}	–	–	–	–	–	–	–	–	–	–	NONE
Ro0.55w	0.55	10^{-4}	–	–	–	–	–	–	–	–	–	–	NONE
Ro0.35w	0.35	10^{-4}	–	–	–	–	–	–	–	–	–	–	NONE

Table 2. Same as Table 1 but for the metastable solutions.

Name	Ro	Λ_i	q	Λ	E_b/E_k	$B_{\text{tor}}^{m=0}$ [10^{14} G]	$B_{\text{pol}}^{m=0}$ [10^{14} G]	B_{dip} [10^{14} G]	$B_s B_\phi / 4\pi$ [10^{27} G 2]	$B_r B_\phi / 4\pi$ [10^{27} G 2]	$B_s^{m=0} B_\phi^{m=0} / 4\pi$ [10^{27} G 2]	$B_r^{m=0} B_\phi^{m=0} / 4\pi$ [10^{27} G 2]	Dynamo
Ro1as	1	$\Lambda(\text{Ro0.75as})$	0.45	70.52	1.18	9.71	0.68	0.47	–	–	–	–	HEM
Ro0.75as	0.75	10	0.064	50.94	1.77	8.13	0.59	0.51	–	–	–	–	DIP
Ro0.75w	0.75	10^{-4}	0.35	0.32	0.01	0.50	0.03	0.08	–	–	–	–	HEM
Ro0.65w	0.65	10^{-4}	0.27	0.36	0.01	0.44	0.03	–	–	–	–	–	HEM
Ro0.25s	0.25	$\Lambda(\text{Ro0.5s})$	0.012	13.60	10.90	4.28	0.26	0.17	4.15	3.59	0.70	1.36	DIP




Figure 1. Time-averaged rms force spectra for the dipolar (left) and hemispherical (right) dynamos at $Ro = 0.75$. The rms forces are averaged over the whole computational domain without excluding boundary layers.

3 FORCE BALANCE

Fig. 1 shows a spectrum of the rms forces in the saturated state of the two Tayler-Spruit dynamo branches, following the formalism of Aubert et al. (2017); Schwaiger et al. (2019). The dipolar dynamo saturates due to a balance between the Lorentz force (red line) and the ageostrophic Coriolis force (dashed green line) at all scales (spectrum on the left in Fig. 1). This confirms the magnetostrophic balance we deduced from Fig. 2 in the Letter. For the hemispherical dynamo, the same balance is found at small scales ($\ell \gtrsim 20$), but at large scales the inertial force is strong enough to be in balance with the ageostrophic Coriolis force and the Lorentz force.

4 TIME-AVERAGED SPECTRA

A wide range of modes ℓ are present in the typical spectra of a dipolar Tayler-Spruit dynamo of Fig. 2. The magnetic spectrum shows the presence of a significant large-scale axisymmetric poloidal field. The even (odd) degrees ℓ dominate in the poloidal (toroidal) axisymmetric magnetic spectra, which confirms the dipolar equatorial symmetry at large scales.

The non-axisymmetric modes triggered differ depending on whether the dynamo is Tayler-instability driven, as seen in Figures 2 and 3. As expected for the Tayler-Spruit dynamo, the dominating mode is the axisymmetric toroidal magnetic field, but we also observe a dominant $m = 1$ mode in the poloidal magnetic energy, which is a signature of the Tayler instability. By contrast, we see that a wider range of orders $m \in [1, 5]$ are present in the poloidal magnetic energy of the kinematic dynamo.

5 MEASURE OF THE SHEAR RATE

The differential rotation is characterized by a dimensionless shear rate $q = r\partial_r \ln \Omega$. We define an effective shear rate based on the time average of the rotation profile in the saturated state. Since it is approximately cylindrical (see Fig. 3), we measure Ω locally at a given height $z = 0.45r_o$ and consider its variation as a function of the cylindrical radius s (see Fig. 4). This allows us to avoid the Ekman layers, which form around the inner shell. In most of the simulations, the shear is found in a broad region centred on the tangent cylinder, especially in $s \in [0.1, 0.5]r_o$. We therefore measure the average slope of the profile in this interval (see Fig. 4):

$$q \equiv \frac{\log \Omega(s = 0.5r_o) - \log \Omega(s = 0.1r_o)}{\log 0.5r_o - \log 0.1r_o} \quad (6)$$

REFERENCES

- Aubert J., Gastine T., Fournier A., 2017, *Journal of Fluid Mechanics*, 813, 558
 Schwaiger T., Gastine T., Aubert J., 2019, *Geophysical Journal International*, 219, S101

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.




Figure 2. Time averaged kinetic (top) and magnetic (bottom) energy density spectra of the dipolar Taylor-Spruit dynamo at $Ro = 0.75$.




Figure 3. Time averaged m -spectra of the magnetic energy density for the hemispherical Taylor-Spruit dynamo at $Ro = 0.85$ (left) and the kinematic dynamo at $Ro = 0.75$ (right).




Figure 4. Rotation profile $\Omega(s)$ at $z = 0.45r_o$ in the simulation $\text{Ro}0.75s$. The green region $s \in [0.1, 0.5] r_o$ is the zone where we measure the effective shear rate q (slope of the blue dashed line). In this example, $q \sim 0.06$. The vertical red line indicates the position of the tangent cylinder.