

Modeling the Solar Wind During Different Phases of the Last Solar Cycle

ZHENGUANG HUANG,¹ GÁBOR TÓTH,¹ NISHTHA SACHDEVA,¹ LULU ZHAO,¹ BART VAN DER HOLST,¹
IGOR SOKOLOV,¹ WARD B. MANCHESTER,¹ AND TAMAS I. GOMBOSI¹

¹*Climate and Space Sciences and Engineering, University of Michigan, Ann Arbor, MI 48109, USA*

ABSTRACT

We describe our first attempt to systematically simulate the solar wind during different phases of the last solar cycle with the Alfvén Wave Solar atmosphere Model (AWSoM) developed at the University of Michigan. Key to this study is the determination of the optimal values of one of the most important input parameters of the model, the Poynting flux parameter, which prescribes the energy flux passing through the chromospheric boundary of the model in the form of Alfvén wave turbulence. It is found that the optimal value of the Poynting flux parameter is correlated with: 1) the open magnetic flux with the Pearson correlation coefficient of 0.913; 2) the area of the open magnetic field regions with the Pearson correlation coefficient of 0.946. These large Pearson correlations suggest that the Poynting flux may vary during a solar cycle, which needs to be investigated with observations, and can shed light on how Alfvén wave turbulence accelerates the solar wind during different phases of the solar cycle. Our results can also be used to set the Poynting flux parameter for real-time solar wind simulations with AWSoM.

1. INTRODUCTION

The solar wind is a continuous plasma flow expanding from the solar corona and propagating through the heliosphere at supersonic speeds as first proposed by [Parker \(1958\)](#). Since the time of its prediction, modeling the solar wind has become an important topic. Over the past few decades, various analytical and numerical magnetohydrodynamic (MHD) models of the solar corona have been developed and successfully applied to simulate the background solar wind (e.g. [Mikić et al. 1999](#); [Groth et al. 2000](#); [Roussev et al. 2003](#); [Cohen et al. 2007](#); [Feng et al. 2011](#); [Evans et al. 2012](#)). Many first-principles models consider Alfvén wave turbulence as the energy source to heat the solar corona and accelerate the solar wind, beginning with early 1D models developed by [Belcher & Davis \(1971\)](#) and [Alazraki & Couturier \(1971\)](#), to 2D models proposed by [Bravo & Stewart \(1997\)](#); [Ruderman et al. \(1998\)](#); [Usmanov et al. \(2000\)](#), and more recently, 3D models including [Lionello et al. \(2009\)](#); [Downs et al. \(2010\)](#); [van der Holst et al. \(2010\)](#). Many physical processes associated with the Alfvén wave turbulence, such as non-linear interactions between forward propagating and reflected Alfvén waves, are included to improve the description of coronal heating ([Velli et al. 1989](#); [Zank et al. 1996](#); [Matthaeus et al. 1999](#); [Suzuki & Inutsuka 2006](#); [Verdini & Velli 2007](#); [Cranmer 2010](#); [Chandran et al. 2011](#); [Matsumoto & Suzuki 2012](#)). Moreover, heat conduction, radiative losses and energy partitioning among particle species as well as temperature anisotropy were introduced in extended MHD (XMHD) models ([Leer & Axford 1972](#); [Chandran et al. 2011](#); [Vásquez et al. 2003](#); [Li et al. 2004](#); [Sokolov et al. 2013](#); [van der Holst et al. 2014](#)). The latest generation of these models is capable of predicting a variety of solar wind observables, including the solar wind density, velocity, the electron and proton (parallel and perpendicular) temperatures, the turbulent wave amplitudes, as well as the wave reflection and dissipation rates, at various locations in the heliosphere.

The Alfvén Wave Solar atmosphere Model (AWSoM) is one of the commonly used first principles Alfvén wave turbulence models developed at the University of Michigan over more than a decade ([van der Holst et al. 2010](#); [Sokolov et al. 2013](#); [Oran et al. 2013](#); [van der Holst et al. 2014](#); [Sokolov et al. 2021](#)). The model has been extensively validated against observations for solar minimum ([Jin et al. 2012](#); [Sachdeva et al. 2019](#)) and maximum conditions ([Sachdeva et al. 2021](#)) includ-

ing comparisons with recent Parker Solar Probe (PSP) encounters (van der Holst et al. 2019, 2022). For these periods of varying solar magnetic activity, the simulated results have been compared to a comprehensive set of observations spanning the low corona to the inner heliosphere.

The Poynting flux parameter (Poynting flux per B ratio) is one of the important inputs for AWSoM, which describes how much Alfvén wave energy is entering into the system to heat the corona and power the solar wind into the inner heliosphere. Sokolov et al. (2013) and van der Holst et al. (2014) estimated this parameter to be approximately $1.1 \text{ MWm}^{-2}\text{T}^{-1}$ based on the chromospheric turbulence observed by *Hinode* (De Pontieu et al. 2007). However the value was modified to $1 \text{ MWm}^{-2}\text{T}^{-1}$ for solar minimum conditions (Sachdeva et al. 2019) and $0.5 \text{ MWm}^{-2}\text{T}^{-1}$ for solar maximum (Sachdeva et al. 2021) conditions to obtain the best agreement with both in-situ and remote observations. It is still unclear how input parameters need to be adjusted to best simulate the solar wind properties for a specific Carrington rotation. This manuscript aims to fill the gap by determining the optimal value of one of the important input parameters of the model, the Poynting flux parameter, during different phases of the solar cycle 24. We will also examine the correlation between the optimal value and the underlying physical quantities/processes.

2. METHODOLOGY

The detailed description of AWSoM can be found in previous publications (Sokolov et al. 2013; Oran et al. 2013; van der Holst et al. 2014; Sokolov et al. 2021). Here we only provide a brief overview. AWSoM is implemented in the BATS-R-US (Block Adaptive Tree Solar Wind Roe-type Upwind Scheme) code (Groth et al. 2000; Powell et al. 1999) within the Space Weather Modeling Framework (SWMF) (Tóth et al. 2005, 2012; Gombosi et al. 2021). The model is driven by the observed radial magnetic field component at the inner boundary located in the lower transition region with a uniform number density ($2 \times 10^{17} \text{ m}^{-3}$) and temperature (50,000 K) distribution. The underlying assumption is that the Alfvén wave turbulence, its pressure and nonlinear dissipation, is the only momentum and energy source for heating the coronal plasma and driving the solar wind, without considering other potential wave heating mechanisms or contributions from small scale reconnections.

Floating boundary condition is applied at the outer boundary so that the simulated solar wind can freely leave the simulation domain.

AWSoM has very few adjustable input parameters. The two important input parameters are the Poynting flux parameter, which is specified as the ratio of the Poynting flux and the magnetic field magnitude at the inner boundary, and the correlation length of the Alfvén wave dissipation (see [van der Holst et al. \(2014\)](#)). The Poynting flux parameter determines the energy input to heat the solar corona and accelerate the solar wind, while the correlation length describes how Alfvén wave turbulence dissipates energy in the solar corona and heliosphere. In this manuscript, we focus on the Poynting flux parameter, which is specified at the inner boundary of AWSoM.

In this study, we simulate one Carrington rotation per year from 2011 to 2019 (the Carrington Rotations and the corresponding magnetogram times are listed in Table 1), using the Air Force Data Assimilative Photospheric flux Transport (ADAPT) Global Oscillation Network Group (GONG) magnetograms ([Hickmann et al. 2015](#)), which are publicly available on <https://gong.nso.edu/adapt/maps/gong>. ADAPT maps use a flux transport model to estimate the radial magnetic field in the regions where there are limited or no observations. There are 12 realizations for each ADAPT map corresponding to 12 different specifications of the supergranulation transport parameters. We randomly picked the seventh realization for all rotations in this manuscript to reduce the cost of computation. In each Carrington rotation, we vary the Poynting flux parameter to obtain different solar wind solutions and compare the simulated solar wind with the OMNI hourly solar wind observations. We then calculate the distance between the simulation results and observations following the methodology introduced by [Sachdeva et al. \(2019\)](#), which quantifies the differences between the simulations and in situ observations at 1 AU to evaluate the performance of the model. The optimal value of the Poynting flux parameter is chosen when the simulated solar wind density and velocity are best compared with the observed values, as these two quantities are most important affecting the CME propagation.

3. SIMULATION RESULTS

Carrington Rotation	UTC Time of the Magnetogram
2106	2011-2-2 02:00:00
2123	2012-5-16 20:00:00
2137	2013-5-28 20:00:00
2154	2014-9-2 20:00:00
2167	2015-8-23 02:00:00
2174	2016-3-3 02:00:00
2198	2017-12-17 02:00:00
2209	2018-10-13 06:00:00
2222	2019-10-2 02:00:00

Table 1. All the ADAPT-GONG magnetograms used in this study

Figure 1 shows the simulated solar wind for two Carrington rotations, one in 2011 (CR2106) near solar minimum and the other in 2013 (CR2137) near solar maximum. Each blue line represents one AWSoM simulation result with a given Poynting flux parameter (while all other parameters are kept the same). The red lines highlight the best run in the corresponding rotation, based on the the best comparison with the observed solar wind density and velocity. The plots illustrate that different values of the Poynting flux parameter can drastically change the simulated solar wind at 1 AU for an active Sun (in 2013). Many simulations give unreasonable results, e.g. very large magnetic field (>25 nT) or solar wind velocity far away from observations. Close to the solar minimum in 2011, the Poynting flux parameter has much less impact, but it is still causes significant variations. In both cases, it is critical to use the correct parameter, otherwise the simulation results will be incorrect.

The differences between the simulated and observed solar wind are quantified by the distances proposed by Sachdeva et al. (2019) and were used to determine the best comparison with observation. We calculate the distances between the simulated and observed densities as well as velocities, as these two quantities significantly impact the CME propagation. We also calculate the average of the density and velocity distances to describe the overall performance. Figure 2, shows that the optimal value of the Poynting flux parameter depends on the choice of the error criteria (for example, based on




Figure 1. AWsoM simulated solar wind velocity, density, temperature and magnetic field (from top to bottom) compared with hourly OMNI observations (black lines) at 1 AU, for the Carrington rotation 2106 (the left panels, a relatively quiet Sun) and 2137 (the right panels, a more active Sun). The blue lines correspond to the simulation results with different Poynting flux parameters in the range of $[0.3, 1.2] \text{ MWm}^{-2}\text{T}^{-1}$ for CR2106, while for CR2137, the range is $[0.15, 1.2] \text{ MWm}^{-2}\text{T}^{-1}$. The red line highlights the results obtained with the optimal value based on the best match with the observed solar wind density and velocity. Note the greater variation in plasma quantities for CR2137.

the density or velocity distances). In this study, we select the optimal value of the Poynting flux parameter when the average distance of the density and velocity reaches the minimum value. Figure 2 shows a monotonic increase of the distance when the Poynting flux parameter is smaller or larger than the optimal value, which means that the optimum is reliably defined. For CR2137 (near solar max), the simulation results become unrealistic when the Poynting flux parameter is larger than $0.75 \text{ MWm}^{-2}\text{T}^{-1}$ as the distances are very large, which confirms that it is critical to choose a correct the Poynting flux parameter in order to obtain reasonable results.

Based on the results shown in these two figures, a natural question is whether we can predict the optimal Poynting flux parameter without performing dozens of simulations. The magnetic field




Figure 2. The distances between the simulated and observed solar wind for CR2106 (the left panel) and CR2137 (the right panel). The x-axis shows the value of the Poynting flux parameter while the y-axis plots the distances between the simulations and OMNI observations. The top row shows the distances between the simulated and observed densities, while the second row shows the distances between the velocities. The third row displays the average of velocity and density distances. The optimal Poynting flux parameter (colored with red) for each panel is found at the minimum of the curves.

structure of the solar corona, for example, may contain some clues. To answer this question, we explore the relationship between the optimal Poynting flux parameter and quantities associated with the magnetic field configurations, including the open magnetic flux (the surface integral of the magnitude of the radial component of the magnetic field $|B_r|$ in the open field regions at the inner boundary), the average magnetic field magnitude on the whole solar surface or in the open field regions, and the area of the open magnetic field regions. We find that the optimal Poynting flux parameter is highly correlated with both the open magnetic flux (shown in Figure 3) with 0.913 Pearson correlation coefficient and the the area of open field regions (see Figure 4) with 0.946 Pearson

correlation coefficient. Figures 3 and 4 show that the optimal Poynting flux parameter, the open magnetic flux and the area of the open field regions are anti-correlated with the Sun’s activity: the values are small during the solar maximum (around 2013-2014) and then increase towards the solar minimum. We performed a linear regression between the optimal Poynting flux parameter P [$\text{MWm}^{-2}\text{T}^{-1}$] and the open magnetic flux F [GR_s^2] as well as the open field area A [R_s^2], and obtained the following following formulas:

$$P = 0.0419 \cdot F - 0.0125 \pm 0.137 \quad (1)$$

$$P = 0.201 \cdot A + 0.0955 \pm 0.109 \quad (2)$$

The \pm terms indicate the standard error of the linear regression.




Figure 3. The left panel shows the open magnetic flux (black, left axis) and the optimal Poynting flux parameter (red, right axis) as a function of the times of the Carrington rotations. The right panel shows the Pearson correlation coefficient r and the linear regression between the open magnetic flux and the optimal Poynting flux parameter with error bars indicating the standard error.

4. SUMMARY AND DISCUSSIONS

Solar wind models based on first principles often assume that Alfvén wave turbulence is the primary energy source to heat the solar corona and accelerate the solar wind. All first principles models need




Figure 4. The left panel shows the open magnetic field area (black, left axis) and the optimal Poynting flux parameter (red, right axis) as a function of the times of the Carrington rotations. The right panel shows the Pearson correlation coefficient r and the linear regression between the area of the open field regions and the optimal Poynting flux parameter with error bars indicating the standard error.

input parameters, which are based on either theoretical expectation or observations. It is important to understand what the physical implications of the input parameters are and how these input parameters would need to be adjusted under different solar conditions, to better understand how the solar corona is heated and the solar wind is accelerated during a full solar cycle, especially if the theory could self consistently explain the acceleration mechanism. In this study, we use AWSoM, which is based on the Alfvén wave turbulence theory, to simulate the solar wind background during different phases of the last solar cycle, and explore how the input parameters need to be adjusted for different solar conditions.

We found that the optimal Poynting flux parameter, which is determined by minimizing the difference between the simulated and observed (by OMNI) solar wind densities and velocities, is highly correlated with the magnetic field structure of the solar corona. To be specific, the open magnetic flux and the area of the open field regions are well correlated with the optimal value of the Poynting flux parameter. The solar cycle dependence of the open magnetic flux and area of the open field regions found in our study are consistent with [Nikolić & \(2019\)](#) and [Lowder et al. \(2017\)](#). On the other hand,

the variation of the optimal Poynting flux parameter, which is defined as the ratio of the Poynting flux and the magnetic field magnitude at the inner boundary, is a new result. Prior work assumed a constant value around $1.1 \text{ MWm}^{-2}\text{T}^{-1}$ (Sokolov et al. 2013; van der Holst et al. 2014), based on the chromospheric turbulence observed by *Hinode* (De Pontieu et al. 2007). However, De Pontieu et al. (2007) is a single observation and we are not aware of any study of the chromospheric turbulence during different phases of the solar cycle. Our study predicts that the chromospheric turbulence may vary during the solar cycle. It will be interesting to see if observations confirm (or contradict) our predictions. A theoretical explanation of these correlations could significantly improve our understanding how the Alfvén wave turbulence heats the solar corona and accelerates the solar wind in different phases of the solar cycle.

This study is also important for the space weather prediction community. First-principles solar wind models have not been used in a real time solar wind prediction primary due to two reasons: 1. high computational cost; 2. the uncertainty of the input parameters. Nowadays, the rapid development of super computers (e.g., the Frontera system supported by NSF and the NASA supercomputing system Pleiades) makes it possible to use a first principles solar wind model to perform real time solar wind predictions, if the input parameters of the model could be specified correctly. The results presented here prescribe one of the important input parameters of AWSoM, the Poynting flux parameter, based on the strong correlation with the open magnetic flux and the area of the open field regions. Both of these quantities can be easily obtained with the required accuracy from the potential field source surface model, for example.

This work was primarily supported by the NSF PRE-EVENTS grant No. 1663800, the NSF SWQU grant No. PHY-2027555, and the NASA Heliophysics DRIVE Science Center (SOLSTICE) at the University of Michigan under grant NASA 80NSSC20K0600. W. Manchester was also supported by the NASA grants NNX16AL12G and 80NSSC17K0686. L. Zhao was also supported by the NASA grants 80NSSC21K0417 and 80NSSC22K0269.

We acknowledge the high-performance computing support from Cheyenne (doi:10.5065/D6RX99HX) provided by NCAR's Computational and Information Systems Laboratory, sponsored by the NSF, and the computation time on Frontera (doi:10.1145/3311790.3396656) sponsored by NSF and the NASA supercomputing system Pleiades.

This work utilizes data produced collaboratively between AFRL/ADAPT and NSO/NISP.

REFERENCES

- Alazraki, G., & Couturier, P. 1971, *A&A*, 13, 380
- Belcher, J. W., & Davis, Leverett, J. 1971, *J. Geophys. Res.*, 76, 3534
- Bravo, S., & Stewart, G. A. 1997, *ApJ*, 489, 992
- Chandran, B. D. G., Dennis, T. J., Quataert, E., & Bale, S. D. 2011, *ApJ*, 743, 197
- Cohen, O., Sokolov, I. V., Roussev, I. I., et al. 2007, *ApJL*, 654, L163
- Cranmer, S. R. 2010, *Astrophys. J.*, 710, 676
- De Pontieu, B., McIntosh, S. W., Carlsson, M., et al. 2007, *Science*, 318, 1574
- Downs, C., Roussev, I. I., van der Holst, B., et al. 2010, *ApJ*, 712, 1219
- Evans, R. M., Opher, M., Oran, R., et al. 2012, *ApJ*, 756, 155
- Feng, X., Zhang, S., Xiang, C., et al. 2011, *ApJ*, 734, 50
- Gombosi, T. I., Chen, Y., Glocer, A., et al. 2021, *Journal of Space Weather and Space Climate*, 11, 42
- Groth, C. P. T., De Zeeuw, D. L., Gombosi, T. I., & Powell, K. G. 2000, *J. Geophys. Res.*, 105, 25053
- Hickmann, K. S., Godinez, H. C., Henney, C. J., & Arge, C. N. 2015, *SoPh*, 290, 1105
- Jin, M., Manchester, W. B., van der Holst, B., et al. 2012, *ApJ*, 745, 6
- Leer, E., & Axford, W. I. 1972, *SoPh*, 23, 238
- Li, B., Li, X., Hu, Y.-Q., & Habbal, S. R. 2004, *Journal of Geophysical Research (Space Physics)*, 109, A07103
- Lionello, R., Linker, J. A., & Mikić, Z. 2009, *ApJ*, 690, 902

- Lowder, C., Qiu, J., & Leamon, R. 2017, *SoPh*, 292, 18
- Matsumoto, T., & Suzuki, T. K. 2012, *ApJ*, 749, 8
- Matthaeus, W. H., Zank, G. P., Oughton, S., Mullan, D. J., & Dmitruk, P. 1999, *ApJL*, 523, L93
- Mikić, Z., Linker, J. A., Schnack, D. D., Lionello, R., & Tarditi, A. 1999, *Physics of Plasmas*, 6, 2217
- Nikolić, & , L. 2019, *Space Weather*, 17, 1293
- Oran, R., van der Holst, B., Landi, E., et al. 2013, *ApJ*, 778, 176
- Parker, E. N. 1958, *ApJ*, 128, 664
- Powell, K. G., Roe, P. L., Linde, T. J., Gombosi, T. I., & de Zeeuw, D. L. 1999, *Journal of Computational Physics*, 154, 284
- Roussev, I. I., Gombosi, T. I., Sokolov, I. V., et al. 2003, *ApJL*, 595, L57
- Ruderman, M. S., Nakariakov, V. M., & Roberts, B. 1998, *A&A*, 338, 1118
- Sachdeva, N., van der Holst, B., Manchester, W. B., et al. 2019, *ApJ*, 887, 83
- Sachdeva, N., Tóth, G., Manchester, W. B., et al. 2021, *Earth and Space Science Open Archive*, 13
- Sokolov, I. V., van der Holst, B., Oran, R., et al. 2013, *ApJ*, 764, 23
- Sokolov, I. V., Holst, B. v. d., Manchester, W. B., et al. 2021, *ApJ*, 908, 172
- Suzuki, T. K., & Inutsuka, S.-I. 2006, *Journal of Geophysical Research (Space Physics)*, 111, A06101
- Tóth, G., Sokolov, I. V., Gombosi, T. I., et al. 2005, *Journal of Geophysical Research (Space Physics)*, 110, 12226
- Tóth, G., van der Holst, B., Sokolov, I. V., et al. 2012, *Journal of Computational Physics*, 231, 870
- Usmanov, A. V., Goldstein, M. L., Besser, B. P., & Fritzer, J. M. 2000, *J. Geophys. Res.*, 105, 12,675
- van der Holst, B., Manchester, W. B., I., Frazin, R. A., et al. 2010, *ApJ*, 725, 1373
- van der Holst, B., Manchester, W. B., I., Klein, K. G., & Kasper, J. C. 2019, *ApJL*, 872, L18
- van der Holst, B., Sokolov, I. V., Meng, X., et al. 2014, *ApJ*, 782, 81
- van der Holst, B., Huang, J., Sachdeva, N., et al. 2022, *ApJ*, 925, 146
- Vásquez, A. M., van Ballegoijen, A. A., & Raymond, J. C. 2003, *ApJ*, 598, 1361
- Velli, M., Grappin, R., & Mangeney, A. 1989, *PhRvL*, 63, 1807
- Verdini, A., & Velli, M. 2007, *ApJ*, 662, 669
- Zank, G. P., Matthaeus, W. H., & Smith, C. W. 1996, *J. Geophys. Res.*, 101, 17093